

Breakup time scale studied in the 8 GeV/c π^- + ^{197}Au reaction

L. Pienkowski ^{a,1}, K. Kwiatkowski ^{b,2}, T. Lefort ^b, W.-c. Hsi ^{b,3},
 L. Beaulieu ^{b,4}, A. Botvina ^c, B. Back ^d, H. Breuer ^e, S. Gushue ^f,
 R.G. Korteling ^g, R. Laforest ^{h,5}, E. Martin ^h,
 E. Ramakrishnan ^h, L.P. Remsberg ^f, D. Rowland ^h,
 A. Ruangma ^h, V.E. Viola ^b, E. Winchester ^h, S.J. Yennello ^h

^a*Heavy Ion Laboratory, Warsaw University, ul. Pasteura 5a, 02-093 Warszawa, Poland*

^b*Department of Chemistry and IUCF, Indiana University, Bloomington, IN 47405, USA*

^c*Institute for Nuclear Research, Russian Academy of Science, 117312 Moscow, Russia*

^d*Physics Division, Argonne National Laboratory, Argonne IL 60439, USA*

^e*Department of Physics, University of Maryland, College Park, MD 20740, USA*

^f*Chemistry Department, Brookhaven National Laboratory, Upton, NY 11973, USA*

^g*Department of Chemistry, Simon Fraser University, Burnaby, BC V5A 1S6, Canada*

^h*Department of Chemistry & Cyclotron Laboratory, Texas A&M University, College Station, TX 77843, USA*

Abstract

Experimental data from the reaction of an 8.0 GeV/c π^- beam incident on a ^{197}Au target have been analyzed in order to investigate the integrated breakup time scale for hot residues. Alpha-particle energy spectra and particle angular distributions supported by a momentum tensor analysis suggest that at large excitation energy, above 3-5 MeV/nucleon, light-charged particles are emitted prior to or at the same time as the emission of the heavy fragments. Comparison with the SMM and GEMINI models is presented. A binary fission-like mechanism fits the experimental data at low excitation energies, but seems unable to reproduce the data at excitation energies above 3-5 MeV/nucleon.

Key words: Light-ion-induced reaction, Multifragmentation, Decay time scale, Thermal excitation energy, Statistical model calculations.

PACS: 25.70.Pq, 21.65.+f, 25.40.-h, 25.80.Hp

¹ E-mail: pienkows@slcj.uw.edu.pl

² Present address: Los Alamos National Laboratory, Los Alamos, NM 87545

³ Present address: 7745 Lake Street, Morton Grove IL 60053

⁴ Present address: Departement de Radio-oncologie, Universitaire de Laval, Quebec, Canada

⁵ Present address: Barnes Hospital, Washington University, St. Louis, MO 63130

During the past decade a major effort has been directed toward studies of the decay of hot nuclei. Experiments with energetic light projectiles offer a unique perspective into decay processes at excitation energies up to about 9 MeV/nucleon. The formation of highly-excited heavy residues in such reactions is less complicated than in heavy-ion reactions, since only a single hot source is formed and small angular momenta and little compression are involved [1,2]. The subsequent evolution of the emission process for heavy fragments as a function of excitation energy is of major concern in understanding the decay mechanisms of hot nuclear matter.

At low excitation energies fission fragments are emitted near the end of the sequential decay chain. At an excitation energy of about 3 MeV/nucleon the emission is delayed by about 10^3 fm/c for the asymmetric split and this delay is much larger for symmetric division [3]. Recently it has been shown that even at E^*/A of about 4 MeV/nucleon, fission fragments from the decay of hot gold-like nuclei are observed with the probability of about 20-40% [4]. The onset of different decay processes involving the prompt emission of multiple fragments is predicted to occur at excitation energies above 4-5 MeV/nucleon [5,6].

The experimental evidence for the transition from a sequential binary decay picture to a simultaneous breakup mechanism is discussed in this letter. Some other signals of the expected transition have recently been presented [7,8] on the basis of the experimental data set analyzed for the purpose of this letter. The Statistical Multifragmentation Model (SMM) [5,9] is one of the “explosive” models and the predictions from this model are compared with the experimental data. A freeze-out volume of $V = 3 \cdot V_0$ (radius $R = 1.44 \cdot R_0$), without any extra expansion energy [8], which describes the $Z \geq 6$ multiplicity and charge distributions at large excitation energies, was assumed for all SMM calculations. Results from the sequential decay model GEMINI [10], including decay time scales and modified to calculate the trajectories in a mutual Coulomb field [11] are also presented. As input for all simulations, the mass and charge of the hot residual nucleus at a given excitation energy were based on event reconstruction procedure of the experimental data [12].

Experiment E900a was performed at the AGS Accelerator at Brookhaven National Laboratory. A secondary 8 GeV/c beam of tagged π^- irradiated a 2 mg/cm² gold target. The ISiS 4π detector array detects charged particles and fragments with charge up to $Z \approx 16$ units and $E/A \approx 1 - 92$ MeV [13]. The experimental methods and previous results can be found in [12,14]. In this letter attention is concentrated on events that detect at least one heavy Intermediate Mass Fragment, IMF, specifically fragments with $Z_{IMF} = 8 - 16$ charge units.

The probability for events with heavy IMFs is presented as a function of E^*/A

in Fig. 1. It is observed that the probability is very low at low E^*/A , increases noticeably in the excitation energy range 3-6 MeV/nucleon and reaches about 40% at about $E^*/A \approx 7-8$ MeV/nucleon. Taking into account 74% detection efficiency, it can be concluded that the selected events are typical events at the excitation energy of about 6-8 MeV/nucleon, where the detected average number of thermal charged particles is about 12-15, the number of fast cascade particle is 10-15, based on a sample of 10^5 events in this E^*/A range. The thermal particles and fragments were selected by imposing a cutoff energy that accepted only fragments below 30 MeV for $Z = 1$ and $9 \cdot Z + 40$ MeV for heavier fragments [15]. SMM and GEMINI models predict for selected events that up to $E^*/A=7-9$ MeV/nucleon the heavy fragment multiplicity with charge $Z \geq 8$ is two or slightly larger than two. The second, mainly heavier, fragment than that detected is too slow to be efficiently detected by ISiS. The event reconstruction procedure gives an estimation of the total missing charge that is consistent with a single missing heavy fragment which is equivalent to an IMF for $E^*/A > 7$ MeV/nucleon. It is observed on Fig. 1 that experimental data are in good agreement with the SMM model predictions up to $E^*/A \approx 8$ MeV/nucleon. The GEMINI model with a density parameter $a=A/10$ MeV $^{-1}$ underestimates the probability to detect heavy IMFs above excitation energies 3-5 MeV/nucleon.

In the 8.0 GeV/c $\pi^- + ^{197}\text{Au}$ reaction the center-of-mass system, CM, is very close to the laboratory system, LAB, $v_{source} \approx 0.01c$. It is observed that the thermal particles and fragments are emitted almost isotropically [16–18], consistent with emission from a thermal-like system. The data were also checked to insure that fast particles were not correlated with the heavy fragments and their spectra were independent of the angle relative to the heavy fragment axis. As shown in model calculations [1,2], the initial spin of the hot nuclei is low, so for the studies of the thermal decay of the hot nuclei the beam axis doesn't define any special direction. Therefore, the direction of the heaviest detected IMF in the LAB system was used as the reference axis for further studies.

Alpha-particle kinetic energy spectra are presented in Fig. 2, for a few selected angles, Θ_{rel} , relative to the axis of the heaviest detected fragment $Z_{IMF} = 8-16$. On the left side of Fig. 2 the spectra at $E^*/A=1-3$ MeV/nucleon are presented. Over most of the angular range the shape of the alpha-particle energy spectra doesn't change much. However, the thermal alpha-particle average kinetic energy, $\langle E_{kin} \rangle$, has a maximum value at small- and large-angles, $\langle E_{kin} \rangle \approx 23$ MeV. The minimum is observed at 50 deg, $\langle E_{kin} \rangle \approx 20$ MeV. The GEMINI model and SMM model (running at $E^*/A \leq 3$ MeV/nucleon essentially as a sequential evaporation model) fit the shape of the spectra, however, predict a more pronounced evolution than the experimental data.

The spectra at $E^*/A=5-7$ MeV/nucleon are presented on the right side of

the Fig. 2. It is important to note that the multifragmentation model SMM reproduces the slope of the spectra up to about 60 MeV. The GEMINI model predicts much steeper spectra. It is also observed that at angles close to the axis defined by a heavy fragment, the alpha-particle kinetic energy spectra have a different shape than the spectra at the larger angles. The average kinetic energy is $\langle E_{kin} \rangle \approx 29$ MeV, compared to a nearly constant value of $\langle E_{kin} \rangle \approx 24$ MeV at the angles between 50 and 170 deg. Similar evolution of the energy spectra is also observed for protons and lithium isotopes, i.e. at large excitation and small angles, Θ_{rel} , the kinetic energy spectra are shifted towards higher energy. The modified GEMINI model [11] predicts a shift similar to the observed one. Tracing the calculations, it was found that at $E^*/A=5-7$ MeV/nucleon GEMINI predicts a short, exponentially distributed time for heavy fragment separation, on average about 100 fm/c. According to the model, most of the alpha-particles are emitted from the hot fragments. The modified version of GEMINI model [11] shows that preferentially low energy alpha-particles are suppressed towards small angles, Θ_{rel} . Particles emitted from the complementary fragment towards the selected fragment are deflected by the Coulomb field to larger angles Θ_{rel} . This result of the model indicates that secondary decay of the hot fragments (emitted promptly after thermalization) is able to explain the kinetic energy shift at small angles, in despite of all doubts concerning the validity of GEMINI model at large excitation energies.

The SMM model doesn't show the observed energy shift of alpha-particles at small angles. This failure seems to be related to the difficulties to define a set of parameters at the freeze-out point configuration. In this context it should be noticed that the version of the SMM model used assumes that secondary decay occurs after the heavy fragments have been fully accelerated, at infinite separation distance, taking into account only two-body Coulomb interaction.

It is interesting to discuss the possibility to describe the shift of alpha-particle kinetic energy at large excitation energy, and at small angles, Θ_{rel} , by a fission-like decay scenario. This requires that the hot nucleus survives as a single self-bound object, the heavy fragments separate after a certain delay time, and the alpha-particles are mainly emitted prior to fragment separation, possibly from a strongly deformed system. One can envision two touching hot spherical nuclei at rest, which would resemble a largely deformed system. In a such a case the alpha-particle energy depends only slightly on the emission angle. Particle emission from the neck at the scission configuration seems to be the only mechanism to enhance the yield of low energy particles emitted perpendicular to the fragments. However, it is observed that the average emitted thermal alpha-particle multiplicity at $E^*/A=5-7$ MeV/nucleon is about 5 and to suppress low energy alpha-particles at small angles, the emission from the neck region would have to be a dominant decay process. In other words, the hot nucleus should emit multiple particles mainly from the strongly de-

formed system, primarily from the neck region, and not from the tip region. Hence, it seems to be difficult to describe such behavior by a fission-like decay mechanism.

The angular multiplicity distributions of thermal particles and light fragments ($Z = 1-7$) relative to the emission axis of a $Z_{IMF} = 8-16$ fragment are shown on the left side of Fig. 3. The particle multiplicity distributions are plotted for selected excitation energy bins, along with SMM model and GEMINI model predictions. In all such angular distribution plots the decrease at angles below 15 deg reflects the ISiS angular granularity. The GEMINI model and SMM model at E^*/A below 3 MeV/nucleon predict a bump at small angles due to particles emitted from the accelerated fragments. The absence of enhanced emission along the heavy IMF axis in the experimental data indicates that the accelerated fragments have lower excitation energies than predicted by the models. Such a picture is consistent with the binary fission scenario, i.e. the fragments are emitted at the end of the sequential decay chain, in a good agreement with refs. [3,4].

The angular distributions at larger E^*/A show that the probability of detecting a particle or fragment close to the selected fragment is suppressed, and suggests that most of the charged particles are emitted almost simultaneous with the selected fragment emission. In other words, the Coulomb field of the heavy fragment makes a cone-shadow not accessible to the other particles emitted at the same time. Consistent with this picture, the SMM model predictions filtered with the detection efficiency procedure describe the experimental data well. At the very large excitation energy, $E^*/A=7-9$ MeV/nucleon, the GEMINI model also fits the data due to the very short time scale predicted by the model: the heavy fragment relative separation time of about 75 fm/c, and the first neutron emission decay time of about 6 fm/c. However, the GEMINI predictions should be taken with the special care at large excitation energies. The predicted decay time at $E^*/A=7-9$ MeV/nucleon is shorter than the time to pass the distance equal to the nucleus diameter with a speed of light. This indicates that the decay time predicted by the GEMINI model is shorter than the energy relaxation time, which contradicts to the model assumption.

In order to investigate systematically the total breakup time of the system and the hypothesis of simultaneous decay, a momentum tensor analysis [19] was performed on the charged particles. The tensor was built for each event from all detected thermal charged particles and light fragments with $Z = 1-7$ charge units. Each element of the tensor, Q_{ij} , was calculated as:

$$Q_{ij} = \sum_{\nu} \frac{p_i^{(\nu)} \cdot p_j^{(\nu)}}{p^{(\nu)}} \quad (1)$$

where, N is the total number of particles in an event, $p^{(\nu)}$ is the particle momentum in the center-of-mass system of N particles with $Z = 1 - 7$ and $p_{i,j}^{(\nu)}$ are the particle momentum Cartesian coordinates. The tensor was diagonalized and the angle between the vector associated with the largest eigenvalue and the emission direction of the selected heaviest fragment, $Z_{IMF} = 8 - 16$, was defined as momentum flow angle, Θ_{flow} . This method permits study of the correlations between the heavy fragments and lighter products of the reaction.

The distributions of the angle Θ_{flow} are presented on the right side of the Fig. 3. The experimental data are almost isotropically distributed at low excitation energy but develop a maximum at 90 deg that becomes increasingly pronounced with increasing excitation energy. An isotropic Θ_{flow} distribution assumption describes more than 95% of events at $E^*/A=1-3$ MeV/nucleon but less than 60% of events at $E^*/A=7-9$ MeV/nucleon. These experimental results should be discussed in a frame of different decay scenarios.

If particles are emitted isotropically, before fragment separation, then Θ_{flow} has also an isotropic distribution. According to the binary fission scenario, this should be the case for the experimental data at E^*/A below 3 MeV/nucleon. The small enhancement of events, below 5%, with Θ_{flow} greater than zero seems to be due to particle emission perpendicular to the fission axis, perhaps from the neck region at the scission configuration.

If there are additional particles emitted from the accelerated fragments, then the momentum ellipsoid should be elongated in the direction of the heavy fragment emission and the Θ_{flow} distribution should be focused at small angles (the kinematical focusing effect). This behavior is shown for low excitation energy by the GEMINI and SMM predictions, corresponding to the angular distributions on the left side of the Fig. 3.

On the other hand, if the associated charged particles are emitted at the same time as the heavy fragment, then preferential Coulomb focusing in a direction perpendicular to the selected heavy fragment emission axis is expected. In such a case the momentum ellipsoid is elongated perpendicular to the heavy fragment emission direction and Θ_{flow} should present a maximum at 90 deg. This focusing effect normal to the heavy fragment axis gives evidence that at high excitation energy the Coulomb focusing effect is stronger than any kinematical focusing effects. The prompt decay scenario, including the Coulomb focusing effect, is contained in the SMM model. Thus, at large excitation energy the SMM model describes the Θ_{flow} distribution well. The modified version of the GEMINI model, with very short emission times after thermalization also fits the Θ_{flow} data at large excitation energy.

The integrated time scale for breakup of the thermal-like system at excitation energies above 5-7 MeV/nucleon can be estimated from the experimental data.

At high excitation energy the observed stronger Coulomb focusing relative to kinematical focusing (right side of the Fig. 3) means that fewer particles are emitted after the acceleration time compared to the number of particles emitted within the first 200 fm/c after fragment formation. The modified GEMINI model gives an estimation that more than about 75% of particles emitted after fragment separation would have to be emitted within the first 200 fm/c to create stronger Coulomb focusing effect than kinematical focusing effect. The light fragment, $Z \leq 7$, angular distributions (Fig. 3) and alpha-particle kinetic energy spectra (Fig. 2) indicate that not many particles can be emitted after the thermalization but before heavy fragment separation. Taking into account these experimental results, it can be concluded that at excitation energies above 5-7 MeV/nucleon the disintegration of the system typically takes place within a total time of approximately 200 fm/c after thermalization. This short lifetime is in a good agreement with the results obtained from the IMF-IMF intensity-interferometry analysis based on the same experimental data [7].

The presented results are obtained for events with detected heavy IMF, $Z = 8 - 16$, and IMF-IMF interferometry analysis was based on the set of events with detected two light IMFs, $Z_{1,2} = 4 - 9$. At a large excitation energies, above 5-7 MeV/nucleon these two set of events represent a large, almost complementary set of events, and cover together more than 70% of detected events. However, it seems that the method based on the sample of events with the splits closer to a symmetric split is a better choice looking for the experimental evidences of the expected transformation from the shape deformation decay mode to the prompt decay mode. Moreover, the set of events with detected heavy fragments covers a large range of excitation energy, the fission-like events were already successfully used in the past [3,20] to study the decay time scale at low excitation energy, and the IMF-IMF interferometry technique has access only to a marginally low fraction of events at low excitation energy due to a very low IMF emission probability.

In conclusion, it is shown that for hot nuclei at excitation energies below 3-5 MeV/nucleon the experimental data are consistent with a sequential binary breakup scenario [3,4], i.e. the heavy fragments are emitted at the end of a sequential decay chain. The fission-like decay scenario above E^*/A of about 6 MeV/nucleon seems unable to fit the experimental data, because it would require emission of too many particles from the neck of the scission configuration. At excitation energies above 3-5 MeV/nucleon, charged-particle emission appears to occur nearly simultaneously with the emission of heavy fragments, $Z_{IMF} = 8 - 16$. The integrated breakup time of the system estimated for the experimental data is approximately 200 fm/c. This observation is in agreement with the delay times from IMF-IMF small angle correlation studies [7] and Statistical Multifragmentation Model (SMM) predictions.

Acknowledgements: This work was supported by the U.S. Department of Energy and National Science Foundation, the National and Engineering Research Council of Canada, Grant No. P03B 048 15 of the Polish State Committee for Scientific Research, Indiana University Office of Research and the University Graduate School, Simon Fraser University and the Robert A. Welch Foundation.

References

- [1] J. Cugnon, Nucl. Phys. A 462 (1987) 751, and J. Cugnon *et al.*, Nucl. Phys. A 470 (1987) 558.
- [2] A.S.Botvina *et al.*, Phys. Lett. B 205 (1988) 421.
- [3] D. Hilscher *et al.*, Phys. Rev. Lett. 62 (1989) 1099.
- [4] U. Jahnke *et al.*, Phys. Rev. Lett. 83 (1999) 4959, and L. Pienkowski *et al.*, Nucl. Phys. A 655 (1999) 269c.
- [5] J.P. Bondorf, A.S. Botvina, A.S. Iljinov, I.N. Mishustin and K. Sneppen, Phys. Rep. 257 (1995) 133.
- [6] D.H.E. Gross, Rep. Prog. Phys. 53 (1990) 605, and D.H.E. Gross, Phys. Rep. 279 (1997) 119.
- [7] L. Beaulieu *et al.*, Phys. Rev. Lett., 84 (2000) 5971.
- [8] T. Lefort *et al.*, Phys. Rev. C 62 (2000) 31604.
- [9] A.S. Botvina, A.S. Iljinov and I.N. Mishustin, Nucl. Phys. A 507 (1990) 649.
- [10] R.J. Charity *et al.*, Nucl. Phys. A 483 (1988) 371, GEMINI-code obtained via ftp from wunmr.wustl.edu.
- [11] W. Gawlikowicz, Acta Phys. Pol. B 28 (1997) 1687.
- [12] L. Beaulieu *et al.*, Phys. Lett. B 463 (1999) 159.
- [13] K. Kwiatkowski *et al.*, Nucl. Instr. Meth. A 360 (1995) 571.
- [14] T. Lefort *et al.*, Phys. Rev. Lett. 83 (1999) 4033.
- [15] K. Kwiatkowski *et al.*, Phys. Lett. B 423 (1998) 21.
- [16] T. Lefort *et al.*, Proceedings of the XXXVII International Winter Meeting on Nuclear Physics, edited by I.Ori Bormio (Italy) 25-30 January 1999, page 7.
- [17] F. Goldenbaum *et al.*, Phys. Rev. Lett. 77 (1996) 1230.
- [18] W.-c. Hsi *et al.*, Phys. Rev. C 60 (1999) 034609.
- [19] J. Cugnon and D. L'Hote, Nucl. Phys. A 397 (1983) 519.
- [20] D. Hilscher and H. Rossner, Ann. Phys. Fr. 17 (1992) 471.




Fig. 1. Dots present the probability to detect an event with a heavy IMF, $Z_{IMF} = 8 - 16$ and solid (dashed) line presents the SMM (GEMINI) model predictions filtered with the experimental detection efficiency. An initial angular momentum of hot nucleus $L = 20\hbar$ was assumed for GEMINI model calculations.




Fig. 2. Alpha-particle energy spectra for selected excitation energy and angular bins relative to the heaviest fragment, $Z_{IMF} = 8 - 16$, emission axis, Θ_{rel} . Solid (dashed) lines present the SMM (GEMINI) prediction filtered with the experimental detection efficiency. The simulation results are normalized to the experimental data assuming the same cross section at a given angle.




Fig. 3. Laboratory angular distributions with respect to the axis defined by the emission of the heaviest detected IMF, $Z_{IMF} = 8 - 16$ charge units, for the selected excitation energy bins. Left side: Average multiplicity of the detected thermal charged particles with $Z = 1 - 7$ charge units as a function of Θ_{rel} . Right side: probability distribution of the momentum flow angle, Θ_{flow} . Solid dots present the experimental data. Solid (dotted) lines present the SMM model predictions filtered (non filtered) with the experimental detection efficiency. Dashed lines are from GEMINI model predictions filtered with the experimental detection efficiency.